



SuperFix

Bactericide / Fungicide



ACTIVE INGREDIENTS

Clove 12.5%, Cottonseed Oil 11.5%, Rosemary 10.6%, Thyme 10.2%, Peppermint Oil 9.5%, Cinnamon 8.2%, Garlic 8.2% -(70.7%)

OTHER INGREDIENTS

White Mineral Oil, Palm Oil, Guar Gum Concentrate, Citrus Pulp, Lemon Juice Concentrate, Citrus Pulp - (29.3%)

TOTAL 100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENT

May cause temporary eye and skin irritation. Do not get in eyes, on skin, or on clothing. Wear protective eyewear such as goggles, face shield, or safety glasses, and chemical resistant gloves. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

FIRST AID

IF IN THE EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. If irritation persists, call a poison control center or doctor for further advice. **IF ON THE SKIN:** Take off contaminated clothing, rinse skin immediately with plenty of soap and water for 15-20 minutes. If irritation persists, call a poison control center or doctor for further advice. **IF SWALLOWED:** Call doctor or poison control center immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by poison control center or doctor. Do not give anything to an unconscious person. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible.

GENERAL INFORMATION

FOR BEST RESULTS SHAKE VIGOROUSLY BEFORE USING

Anjon AG **SuperFix** is a concentrate and is designed for use both in greenhouses and outdoors. Anjon AG **SuperFix** is intended to be mixed with water and applied with pressurized power operated sprayers or delivered through drip irrigation systems. When diluting, it is recommended to add approximately half the water to the applicator tank, then add the proper amount of Anjon AG **SuperFix** in accordance with the rate table, agitate, then add the remaining amount of water. Then agitate the mixture to insure proper suspension. Anjon AG **SuperFix** fungicide pest control agent. Anjon AG **SuperFix** by the product label. A small volume test mix is recommended to ensure compatibility. Anjon AG **SuperFix** may cause a phytotoxic damage to green scale test to ensure lack of phytotoxicity of the combination is recommended.

DIRECTIONS OF USE

Anjon AG **SuperFix** is a multi-purpose broad spectrum bactericide/fungicide. Anjon AG **SuperFix** can be used for treatment of all food crops and ornamentals. Anjon AG **SuperFix** can also be used for soil applications to aid in reducing the damaging effects of soil borne bacterial and fungal diseases.

APPLICATION GUIDELINES

Anjon AG **SuperFix** does not normally require a pH change. It is recommended that a pH in the less be used.

APPLICATION TIMING

Apply diluted spray when high humidity for the development of disease symptoms in crops or at first sign of disease and continue applications at 7-10 day intervals or until disease symptoms are gone. For soil with severe potato scab apply one week prior to planting and then reapply at planting. For reapplication apply at time of planting. Follow with foliar spray applications at 8-10 inches of foliage.

MINIMUM TIME TO APPLICATION TO HARVEST

Spraying or drip application of the product can occur up to the time of harvest on all intended crops.

APPLICATION METHOD AND EQUIPMENT

Anjon AG **SuperFix** can be sprayed on foliage, the surface of soil a drip irrigation system at the rates on the appropriate tables. When sprayed on f hours of a dry period so that irrigation or rain does not wash off application. When applied topically to soil, immediately follow application of Anjon AG **SuperFix** water to wash into soil.

For drip applications apply to soil that is with humidity level of 70% or more. After Anjon AG **SuperFix** has cleared lines follow with enough water to wash down to root level. Always wash into soil after application with plain water to make sure that Anjon AG **SuperFix** penetrates down to the depth of the root ball. Consult spray nozzle and accessory catalogues for specific information on proper equipment calibration. For best coverage use small droplets.

STORAGE AND DISPOSAL

Store in original UV protected containers only. Store at temperatures between 41°F to 85°F (5°C to 30°C). Disposal: Wastes resulting from using this product may be disposed of on site. Dispose of empty jug in a sanitary land fill as allowed by state and local authorities.

CONDITIONS & LIABILITIES

Anjon AG, Inc. (Seller) warrants that the chemical composition of this product conforms to the chemical description given on this label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS EXPRESSED, IMPLIED, OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. No one is authorized to make any other warranty, guarantee or directions concerning this product, and no such warranties, guarantees or directions shall be valid or binding upon Seller. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation, are beyond the control of Seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. In no case shall Seller be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition of sale by Seller and is accepted as such by the buyer. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option, as full discharge of Seller's liability.

- NET CONTENTS: 1 Gal (3.79 L) Jug
- Jug NET CONTENTS: 2.5 Gal (9.46 L) Jug
- Jug NET CONTENTS: _____

Specific Crops	Target Diseases	App. Rate fl oz	Rate per Acre Gallons	Application Directions & Interval
Blueberry	Mummy Berry Anthracnose Fruit Rot Bacteria Soft Rot Botrytis Fruit Rot Phomopsis Leaf Spot, Twig Blight & Fruit Rot Powdery Mildew Rhizopus	12 16 12	30-50	Spray at green tip, early pink bud, pink bud, early bloom, mid bloom, petal fall and fruit set
Cucurbits Cucumber, squash, pumpkin, melon, pickle	Bacterial Leaf Spot Bacteria Wilt Bacteria Fruit Blotch Downey Mildew Powdery Mildew Anthracnose Late Blight Rhizopus	14 14 12 10 10	25-30 25-40 30-40 30-40 25-40	Preventative - make application every 7 days. After first symptoms observed, apply every 3-5 days until control is established. Recommended to use in combination with EF400.
Cotton	Bacteria Blight	16	10-15	Apply in combination with EF400
Dry Edible Beans and Green Beans	Common Blight Halo Blight Brown Spot White Mold (Sclerotinia) Powdery Mildew Downey Mildew	12 12 12	20 20 20	Apply 3rd trifoliolate, R1, R4, R7. Recommended use in combination with EF400 to reduce fungal problems.
Field Corn	Anthracnose Goss's Wilt	14 16	20 20	Anthracnose - Spray at 4th leaf and continue every 10-14 days or 7-10 days under high pressure conditions. For Goss's Wilt apply at V-4 stage and row tassel. Apply every 7-10 days until disease is suppressed from stalk.
Onion	Anthracnose - onion smudge	14	25-40	Begin shortly after emergence and spray every 7 days
Peppers	Bacteria Leaf Spot	12	25-40	If first symptoms are observed apply every 5-7 days until control is attained.
Potatoes	Common Scab Bacteria Wilt Early Blight Late Blight	14 14 14 14	25-30 25-30 25-30 25-30	For soil treatment for Scab use in combination with EF400. After application water into soil to reach root area. For topical spray use as recommended. Begin spraying at first sign of disease.
Potato Seed Preplant	To help stop the spread of Scab and other bacterial and fungal disease	16	20	Use in combination with EF400. Use fine spray and allow to dry before planting. Good coverage is critical.
Soybeans	Bacteria Blight	12	20	Application schedule: R1, R4, R-6
Strawberry	Bacteria Blight Bacteria Wilt Botrytis Powdery Mildew	12 12 12 12	25-40 25-40 25-40 25-40	Product should be applied on a preventative program. For soil borne diseases can applied through drip irrigation system. Use in combination with EF400
Tomatoes	Bacteria Leaf Spot Bacterial Canker Bacteria Wilt Early Blight Late Blight	12 12 12 12	25-40 25-40 25-40 25-40	Begin spraying at first signs of disease onset and maintain a 7-10 day interval during periods of high infection
Wheat	Fusarium Head Blight Bacteria Head Blight	16 16	20 20	4th to 5th leaf followed at 14 days later
Aerial Application	Use a pH of 6.0 to 7.0 and a dilution rate of not less than 2% or 2 gallons per 100 gallons.	2%	5 gal/ acre	Apply at first signs of disease onset. Repeat at 7 to 10 days.

Guaranteed by:



220 ADAMS DR. #280-221, WEATHERFORD, TX 76086 • 855-434-0300